

NEW TYPE SCALAR FIELDS FOR COSMIC ACCELERATION

Alex Kehagias*

*Department of Physics, National Technical University of Athens,
GR-15773, Zografou, Athens, Greece*

Abstract

We present a model where a non-conventional scalar field may act like dark energy leading to cosmic acceleration. The latter is driven by an appropriate field configuration, which result in an effective cosmological constant. The potential role of such a scalar in the cosmological constant problem is also discussed.

*e-mail: kehagias@cern.ch

During the last years, evidence of an accelerating expansion of the Universe, first published in [1],[2], has become one of the most striking results in modern cosmology and in physics, in general. The most obvious cause of this acceleration is a cosmological constant, although a large number of alternatives have been proposed [3]. Here we will discuss a rather simple, although non-conventional, scalar model, which may also account for the present Universe acceleration.

The scalar field we will consider is a real one and its dynamics is determined by the action

$$S = \int d^4x \sqrt{-g} \left(-\frac{1}{2} \partial_\mu \Phi(x) \partial^\mu \Phi(x) + k \partial_\mu \Phi(x) \partial^\mu \Phi(x+\alpha) \right). \quad (1)$$

The parameter k is a numerical constant and the shifts $(x+\alpha)$ in the argument of the scalar field are generated by a 4-vector α^μ , which has only a time component, i.e., $\alpha^\mu = (\alpha, 0, 0, 0)$. Hence, for $x^\mu = (t, x_1, x_2, x_3)$ we have $(x+\alpha) = (t+\alpha, x_1, x_2, x_3)$. We assumed for simplicity that there is no potential for the scalar field. The unusual feature of the action (3) is that, besides the standard kinetic term, there is also an additional non-local derivative term¹. A similar term has been discussed first time to my knowledge, by Feynman in his thesis [4] in a classical mechanical context, and in connection to the theory of action at a distance. It can be considered as describing a field interacting with itself in a distant mirror by means of advanced and retarded waves. In a modern picture, modelling our universe as a 3-brane, it could be the result of a scalar interacting in a distant 3-brane through bulk waves. Alternatively, expanding for $t \gg a$, we see that

$$S = \int d^4x \sqrt{-g} \left(-\left(\frac{1}{2} - k\right) \partial_\mu \Phi(x) \partial^\mu \Phi(x) + k \sum_{n=2} \frac{1}{n!} \partial_\mu \Phi(x) a^{\mu_1} \dots a^{\mu_n} \partial^{\mu_1 \dots \mu_n} \Phi(x) \right), \quad (3)$$

and for $k < 1/2$, we have a higher-derivative theory. Irrespectively of its origin, we would like to couple this scalar field to gravity and find cosmological solutions for the corresponding gravity-scalar system.

The scalar field equation can be obtained from the action (3), after scalar variations $\delta\Phi(x)$, and the resulting equation is

$$\nabla^2 \Phi(x) - k \nabla^2 \Phi(x+\alpha) - k \nabla^2 \Phi(x-\alpha) = 0. \quad (4)$$

Similarly, the induced energy-momentum tensor is given by

$$T_{\mu\nu}^{(\Phi)} = \partial_\mu \Phi(x) \partial_\nu \Phi(x) - k \partial_\mu \Phi(x) \partial_\nu \Phi(x+\alpha) - k \partial_\mu \Phi(x+\alpha) \partial_\nu \Phi(x)$$

¹This modifies the propagator of the scalar field, which for the action (3) is given by

$$\Delta = \left\{ p^2 [1 - 2k \cos(p^\mu \alpha_\mu)] \right\}^{-1}. \quad (2)$$

For $|k| < 1/2$ it has a single pole at $p^2 = 0$.

$$-\frac{1}{2}g_{\mu\nu}[\partial_\rho\Phi(x)\partial^\rho\Phi(x)-2k\partial_\rho\Phi(x)\partial^\rho\Phi(x+\alpha)] , \quad (5)$$

and, according to the standard prescription, gravity should coupled to it. However, this is not possible in the present case and $T_{\mu\nu}^{(\Phi)}$ cannot be the right-hand side of Einstein equations. The reason is that, due to the second term in the action(3), general coordinate invariance is lost and consequently, $T_{\mu\nu}^{(\Phi)}$ is not conserved. Indeed, it can easily be verified that the covariant divergence of $T_{\mu\nu}^{(\Phi)}$ does not vanish. Instead we find that

$$\nabla_\mu T^{(\Phi)\mu}{}_\nu = k\nabla^2\Phi(x-\alpha)\partial_\nu\Phi(x) - k\nabla^2\Phi(x)\partial_\nu\Phi(x+\alpha) . \quad (6)$$

Gravity must couple then to an “improved” energy-momentum tensor $\Theta_{\mu\nu}$, which we define as

$$\Theta_{\mu\nu} = T_{\mu\nu}^{(\Phi)} + t_{\mu\nu} , \quad (7)$$

and it is such that $\nabla_\mu\Theta^{\mu\nu} = 0$. Hence, the extra term $t_{\mu\nu}$ must satisfy

$$\nabla_\mu t^\mu{}_\nu = k\nabla^2\Phi(x)\partial_\nu\Phi(x+\alpha) - k\nabla^2\Phi(x-\alpha)\partial_\nu\Phi(x) , \quad (8)$$

in order $\Theta_{\mu\nu}$ to have vanishing covariant divergence. Einstein equations are then written as

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \kappa^2\Theta_{\mu\nu} , \quad (9)$$

which is now a consistent system to be solved.

We will look now for cosmological solutions to eqs(4,9) of the form

$$ds^2 = -dt^2 + a(t)^2(dx^2 + dy^2 + dz^2) , \quad \Phi = \Phi(t) . \quad (10)$$

Then, we get from eq.(5) that the induced energy-momentum tensor is

$$T^{(\Phi)\mu}{}_\nu = \text{diag}(-\rho_\Phi, p_\Phi, p_\Phi, p_\Phi) \quad (11)$$

where

$$\rho_\Phi = \frac{1}{2}\dot{\Phi}(t)^2 - k\dot{\Phi}(t)\dot{\Phi}(t+\alpha) , \quad p_\Phi = \rho_\Phi . \quad (12)$$

Moreover, defining

$$t^\mu{}_\nu = \text{diag}(-\rho, p, p, p) , \quad (13)$$

we find that $t_{\mu\nu}$ satisfies eq.(8) and $\Theta_{\mu\nu} = T_{\mu\nu}^{(\Phi)} + t_{\mu\nu}$ is conserved for

$$\begin{aligned} \rho &= k \int_t^{t+\alpha} \ddot{\Phi}(\tau-\alpha)\dot{\Phi}(\tau)d\tau , \\ p &= k\dot{\Phi}(t+\alpha)\dot{\Phi}(t) - k\dot{\Phi}(t-\alpha)\dot{\Phi}(t) - k \int_t^{t+\alpha} \ddot{\Phi}(\tau-\alpha)\dot{\Phi}(\tau)d\tau . \end{aligned} \quad (14)$$

Indeed, the conservation of $\Theta_{\mu\nu}$ is equivalent to

$$\dot{\rho}_\Phi + \dot{\rho} + 3\frac{\dot{a}}{a}(\rho_\Phi + \rho + p_\Phi + p) = 0, \quad (15)$$

which can easily be verified for $\rho_\Phi, p_\Phi, \rho, p$ of eqs.(12,14). Then, the improved energy-momentum tensor $\Theta_{\mu\nu}$ has the form

$$\Theta_\nu^\mu = \text{diag}(-\rho_\Theta, p_\Theta, p_\Theta, p_\Theta), \quad (16)$$

where

$$\begin{aligned} \rho_\Theta = \rho_\Phi + \rho &= \frac{1}{2}\dot{\Phi}(t)^2 - k\dot{\Phi}(t+\alpha)\dot{\Phi}(t) + k \int_t^{t+\alpha} \ddot{\Phi}(\tau-\alpha)\dot{\Phi}(\tau)d\tau, \\ p_\Theta = p_\Phi + p &= \frac{1}{2}\dot{\Phi}(t)^2 - k\dot{\Phi}(t-\alpha)\dot{\Phi}(t) - k \int_t^{t+\alpha} \ddot{\Phi}(\tau-\alpha)\dot{\Phi}(\tau)d\tau \end{aligned} \quad (17)$$

and Einstein (9) and scalar equations (4) are explicitly written as

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{\kappa^2}{3}\rho_\Theta, \quad (18)$$

$$-2\frac{\ddot{a}}{a} - \frac{\dot{a}^2}{a^2} = \kappa^2 p_\Theta, \quad (19)$$

$$\begin{aligned} &\ddot{\Phi}(t) - k\ddot{\Phi}(t+\alpha) - k\ddot{\Phi}(t-\alpha) + \\ &+ 3\frac{\dot{a}}{a}(\dot{\Phi}(t) - k\dot{\Phi}(t+\alpha) - k\dot{\Phi}(t-\alpha)) = 0. \end{aligned} \quad (20)$$

A first integral of the scalar equation (20) is

$$\dot{\Phi}(t) - k\dot{\Phi}(t+\alpha) - k\dot{\Phi}(t-\alpha) = c a(t)^{-3}, \quad (21)$$

where c is an integration constant. As usual, only two of the equations (18,19,20) are independent. We take Friedman equation (18) and the integrated scalar equation (21). With a vanishing $c(=0)$, the solution in the usual case $k=0$ is $\Phi = \text{constant}$. However, in the present case where $k \neq 0$, other solutions exist as well. Indeed, one may easily verify that a solution to eq.(21) with $c=0$ is provided by the configuration

$$\Phi(t) = m \left(\cosh(\mu t) + \nu \sinh(\mu t) \right), \quad \mu = \alpha^{-1} \text{arccosh}\left(\frac{1}{2k}\right) \quad (22)$$

where m, μ are mass parameters and ν is a numerical one. Then, the right-hand side of Friedman equation (18) turns out to be

$$\rho_\Theta = \frac{m^2 \sqrt{1-4k^2}}{4a^2} (1-\nu^2) \text{arccosh}\left(\frac{1}{2k}\right)^3, \quad (23)$$

which is a real constant for $0 < k < 1/2$. Moreover, we also get that

$$p_{\Theta} = -\frac{m^2\sqrt{1-4k^2}}{4a^2}(1-\nu^2)\operatorname{arccosh}\left(\frac{1}{2k}\right)^3 = -\rho_{\Theta}, \quad (24)$$

which is the typical equation of state for cosmological constant. Thus, in this model, the scalar leads to an effective cosmological constant Λ_{Φ} with value

$$\Lambda_{\Phi} = \frac{\kappa^2 m^2 \sqrt{1-4k^2}}{4a^2}(1-\nu^2)\operatorname{arccosh}\left(\frac{1}{2k}\right)^3. \quad (25)$$

The parameter μ is of the order of α^{-1} , and the observed cosmic acceleration could be driven by a field like the present one for $m^2 \approx 10^9 \alpha^2 (1-\nu^2)^{-1} \text{cm}^{-4}$. Thus, a scalar field with dynamics described by the action (3) is a candidate for dark energy. Moreover, it can also be considered, in principle, as an inflaton field for an early time inflation.

We should note here that our scalar field may also be used for the cosmological constant problem [5]. In the presence of a bare cosmological constant Λ , Einstein equations (9) are written as

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = \kappa^2 T_{\mu\nu}, \quad (26)$$

where $T_{\mu\nu}$ is the matter energy-momentum tensor. With a non-zero Λ , flat space-time is not a solution, namely, $g_{\mu\nu} = \eta_{\mu\nu}$ does not satisfy (26). However, for $T_{\mu\nu} = \Theta_{\mu\nu}$ the bare cosmological constant Λ is shifted to $\Lambda + \Lambda_{\Phi}$ by the scalar field (22) and it can take any value by appropriate choice of the numerical parameter ν . Then, by adjusting ν so that $\Lambda_{\Phi} = -\Lambda$, flat space-time is now a solution to eq.(26). This is always possible for a negative bare cosmological constant, whereas, for a positive one, the parameter ν should be such that $\nu > 1$. In the latter case, although ρ_{Θ} is negative violating seemingly weak energy conditions, the energy density ρ_{Φ} of the scalar field associated with the canonical energy-momentum tensor is positive for $t > 0$. In other words, for any value of Λ , there always exists a scalar described by the action (3) and profile given by (22), which may nullify Λ or drive it to its present value. The question why this value is so tiny remains open.

Acknowledgement. This work is supported by the EPEAEK programme ‘‘Pythagoras’’ and co-funded by the European Union (75%) and the Hellenic State (25%).

References

- [1] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], *Astrophys. J.* **517**, 565 (1999) [arXiv:astro-ph/9812133].

- [2] A. G. Riess *et al.* [Supernova Search Team Collaboration], *Astron. J.* **116**, 1009 (1998) [arXiv:astro-ph/9805201].
- [3] E. J. Copeland, M. Sami and S. Tsujikawa, arXiv:hep-th/0603057.
- [4] R. P. Feynman, “The Principle Of Least Action In Quantum Mechanics,”
- [5] S. Weinberg, *Rev. Mod. Phys.* **61**, 1 (1989).